arXiv:0801.4723v1 [math.LO] 30 Jan 2008

February 3, 2008

RAMSEY-LIKE EMBEDDINGS

VICTORIA GITMAN

Abstract. One of the numerous equivalent characterizations of a Ramsey cardinal κ involves the existence of certain types of elementary embeddings for transitive sets of size κ satisfying a large fragment of ZFC. I introduce new large cardinal axioms generalizing the Ramsey embeddings and show that they form a natural hierarchy between weakly compact cardinals and measurable cardinals. These new axioms serve to further our knowledge about the elementary embedding properties of smaller large cardinals, in particular those still consistent with $V = L$.

1. Introduction

Most large cardinals, including measurable cardinals and stronger notions, are defined in terms of the existence of elementary embeddings with that cardinal as the critical point. Several smaller large cardinals, such as weakly compact, indescribable, and Ramsey cardinals, have definitions in terms of elementary embeddings, but are more widely known for their other properties (combinatorial, reflecting, etc.). Other smaller large cardinals, such as ineffable and subtle cardinals, have no known elementary embedding properties. I will investigate the elementary embedding property of Ramsey cardinals and introduce new large cardinal axioms by generalizing the Ramsey embeddings. By placing these new large cardinals within the existing hierarchy, I try to shed light on the variety of elementary embedding properties that are possible for smaller large cardinals. The new large cardinals also have elementary embeddings more suitable for use in indestructibility arguments than the Ramsey embeddings. In a forthcoming joint paper with Thomas Johnstone, [**?**], we use the embedding properties to obtain some basic indestructibility results for Ramsey cardinals through the techniques of lifting embeddings. I hope that this project will motivate set theorists who work with smaller large cardinals to focus on investigating their elementary embedding properties. I will begin with some basic definitions and terminology.

Let ZFC$^-$ denote the fragment of ZFC consisting of ZF without the Powerset axiom and a form of Choice which states that every set is bijective with some ordinal. I will call a transitive set $M \models$ ZFC$^-$ of size κ with $\kappa \in M$ a *weak κ-model*. I will say that a weak κ-model M is a *κ-model* if additionally $M^{<\kappa} \subseteq M$. Observe that for any cardinal κ, if $M \prec H_{\kappa^+}$ has size κ with $\kappa \subseteq M$, then M is a weak κ-model. Similarly, if $\lambda > \kappa$ is a regular cardinal and $X \prec H_\lambda$ has size κ with $\kappa + 1 \subseteq X$, then the Mostowski collapse of X is a weak κ-model. So there are always many weak κ-models for any cardinal κ. If additionally $\kappa^{<\kappa} = \kappa$, we can use a Skolem-Lowenheim type construction to build κ-models $M \prec H_{\kappa^+}$ and substructures $X \prec H_\lambda$ whose collapse will be a κ-model. Unless specifically stated otherwise, the sources and targets of elementary embeddings are *assumed to be transitive*. I will call an elementary embedding $j : M \to N$ *κ-powerset preserving* if it has critical point κ and M and N have the *same* subsets of κ. For example, it is trivially true that any elementary embedding $j : V \to M$ with critical point κ such that $M \subseteq V$ is κ-powerset preserving.

I will define the Ramsey-like embedding properties shortly. To provide a motivation for this collection of definitions, I will begin by recalling the various equivalent definitions of weakly compact cardinals.

Theorem 1.1. *If $\kappa^{<\kappa} = \kappa$, then the following are equivalent.*

(1) *κ is weakly compact. That is, κ is uncountable and every κ-satisfiable theory in a $L_{\kappa,\kappa}$ language of size at most κ is satisfiable.*
(2) *For every $A \subseteq \kappa$, there is a transitive structure W properly extending V_κ and $A^* \subseteq W$ such that $\langle V_\kappa, \in, A\rangle \prec \langle W, \in, A^*\rangle$.*
(3) *κ is inaccessible and every κ-tree has a cofinal branch.*
(4) *Every $A \subseteq \kappa$ is contained in weak κ-model M for which there exists an elementary embedding $j : M \to N$ with critical point κ.*
(5) *Every $A \subseteq \kappa$ is contained in a κ-model M for which there exists an elementary embedding $j : M \to N$ with critical point κ.*
(6) *Every $A \subseteq \kappa$ is contained in a κ-model $M \prec H_{\kappa^+}$ for which there exists an elementary embedding $j : M \to N$ with critical point κ.*
(7) *For every κ-model M, there exists an elementary embedding $j : M \to N$ with critical point κ.*
(8) *For every κ-model M, there exists an elementary embedding $j : M \to N$ with critical point κ such that j and M are elements of N.*

For a proof of these equivalences see [**?**] (ch. 6). Now we are ready to state the large cardinal notions that will be the focus of this paper. The general idea is to consider the various equivalent elementary embedding properties of weakly compact cardinals with the added assumption that the embeddings have to be κ-*powerset preserving*. We will soon see that this additional assumption destroys the equivalence in the strongest possible sense.

Definition 1.2 (Weak Ramsey Embedding Property)**.** A cardinal κ has the *weak Ramsey embedding property* if every $A \subseteq \kappa$ is contained in a weak κ-model M for which there exists a κ-powerset preserving elementary embedding $j : M \to N$. We say that a cardinal is *weakly Ramsey* if it has the weak Ramsey embedding property.

Definition 1.3 (Ramsey Embedding Property)**.** A cardinal κ has the *Ramsey embedding property* if every $A \subseteq \kappa$ is contained in a weak κ-model M for which there exists a κ-powerset preserving elementary embedding $j : M \to N$ satisfying the property that whenever $\langle A_n \mid n \in \omega\rangle$ is a sequence of subsets of κ such that each $A_n \in M$ and $\kappa \in j(A_n)$, then $\cap_{n\in\omega} A_n \neq \emptyset$.

For $\langle A_n \mid n \in \omega\rangle \in M$, of course, the conclusion follows trivially. So the content here is for sequences *not* in M. I will show later that a cardinal is Ramsey if and only if it has the Ramsey embedding property (this follows from results in [**?**], [**?**], and [**?**] among others).

Definition 1.4 (Strong Ramsey Embedding Property)**.** A cardinal κ has the *strong Ramsey embedding property* if every $A \subseteq \kappa$ is contained in a κ-model M for which there exists a κ-powerset preserving elementary embedding $j : M \to N$. We say that a cardinal is *strongly Ramsey* if it has the strong Ramsey embedding property.

Definition 1.5 (Super Ramsey Embedding Property)**.** A cardinal κ has the *super Ramsey embedding property* if every $A \subseteq \kappa$ is contained in a κ-model $M \prec H_{\kappa^+}$ for which there exists a κ-powerset preserving elementary embedding $j : M \to N$. We say that a cardinal is *super Ramsey* if it has the super Ramsey embedding property.

Definition 1.6 (Total Ramsey Embedding Property)**.** A cardinal κ has the *total Ramsey embedding property* if for every κ-model $M \prec H_{\kappa^+}$, there exists a κ-powerset preserving elementary embedding $j : M \to N$. We say that a cardinal is *totally Ramsey* if it has the total Ramsey embedding property.

Note that since a weak κ-model can take the Mostowski collapse of any of its elements, the definitions above hold not just for any $A \subseteq \kappa$, but more generally for any $A \in H_{\kappa^+}$. Also, observe that the "total Ramsey embedding property" → "super Ramsey embedding property" → "strong Ramsey embedding property" → "Ramsey embedding property" → "weak Ramsey embedding property".

The following theorem, which will be proved in parts in the subsequent sections, summarizes what is known about the placement of these new large cardinals in the large cardinal hierarchy. Also, see the diagram on the next page.

Theorem 1.7.

(1) *A measurable cardinal is a super Ramsey limit of super Ramsey cardinals.*
(2) *A super Ramsey cardinal is a strongly Ramsey limit of strongly Ramsey cardinals and an ineffably Ramsey limit of ineffably Ramsey cardinals.*
(3) *A strongly Ramsey cardinal is a Ramsey limit of Ramsey cardinals and a limit of ineffably Ramsey cardinals.*
(4) *A cardinal has the Ramsey embedding property if and only if it is Ramsey.*
(5) *A Ramsey cardinal is a weakly Ramsey limit of weakly Ramsey cardinals.*[1]
(6) *Weakly Ramsey cardinals are consistent with* $V = L$.[2]
(7) *A weakly Ramsey cardinal is a weakly ineffable limit of ineffable cardinals, but not necessarily ineffable.*
(8) *There are no totally Ramsey cardinals.*

Theorem 1.7 shows, surprisingly, that the various embedding properties that define weakly compact cardinals form a hierarchy of strength when you add the powerset preservation property, and one such property is even inconsistent!

---

[1] This follows from a result of Ian Sharpe [**?**] to be discussed in Section 5.

[2] I am grateful to Philip Welch for answering this question.

The solid arrows indicate direct implications and the dashed arrows indicate relative consistency.



Definition 1.8. Suppose M is a transitive model of ZFC^- and λ is a cardinal in M. Then $U \subseteq \mathcal{P}(\lambda) \cap M$ is an *M-ultrafilter* if $\langle M, U\rangle \models$ "U is a normal ultrafilter".

Recall that an ultrafilter U is *normal* if every regressive function $f : A \to \lambda$ with $A \in U$ is constant on some $B \in U$. Note that we are taking normality from the point of view of M as part of the definition of an M-ultrafilter.

Proposition 1.9. *Suppose M is a weak κ-model such that $M^\alpha \subseteq M$ for some $\alpha < \kappa$ and $j : M \to N$ is a κ-powerset preserving embedding. Let $X = \{j(f)(\kappa) \mid f : \kappa \to M \text{ and } f \in M\} \subseteq N$ and let $\pi : X \to K$ be the Mostowski collapse. Then $h = \pi \circ j : M \to K$ is an elementary embedding with critical point κ having the properties:*

1. *h is κ-powerset preserving,*
2. *h is an ultrapower by an M-ultrafilter on κ,*
3. *$\kappa \in j(A)$ if and only if $\kappa \in h(A)$ for all $A \subseteq \kappa$ in M,*
4. *K has size κ,*
5. *$K^\alpha \subseteq K$,*
6. *we get the commutative diagram:*

$$\begin{tikzcd}
M \arrow[d, "h"'] \arrow[dr, "j"] & \\
K \arrow[r, "\pi^{-1}"] & N
\end{tikzcd}$$

Proof. Clearly the size of K is κ since M has size κ. Define $U = \{A \subseteq \kappa \mid A \in M \text{ and } \kappa \in j(A)\}$. It follows by standard seed theory arguments that U is an M-ultrafilter, the embedding h is the ultrapower by U, and $K^\alpha \subseteq K$ [?] (ch. 0). For (1), observe that $\pi(\beta) = \beta$ for all $\beta \leq \kappa$, and hence the critical point of π^{-1} is above κ. This implies that N and K have the same subsets of κ. Finally, for (3), observe that $\kappa \in j(A) \leftrightarrow \pi(\kappa) \in \pi \circ j(A) \leftrightarrow \kappa \in \pi \circ j(A) \leftrightarrow \kappa \in h(A)$. $\square$

By Proposition 1.9, we can always assume in the definitions of Ramsey-like embeddings that $j : M \to N$ is an *ultrapower* by an M-ultrafilter on κ, the target N has size κ, and if $M^\alpha \subseteq M$, then $N^\alpha \subseteq N$. Also, from condition (3), it follows that the Ramsey embedding property can be restated in terms of having a κ-powerset preserving elementary embedding $j : M \to N$ by an M-ultrafilter U on κ, with the property that whenever $\langle A_n \mid n \in \omega\rangle$ is a sequence of elements of U, then $\cap_{n\in\omega} A_n \neq \emptyset$.

2. The Large Cardinal Hierarchy

In this section, I show that the definitions above do define large cardinal notions, and discuss where these large cardinals fit into the large cardinal hierarchy. The arguments below all rely on the powerful reflecting properties of Ramsey-like embeddings, which arise from the fact that the source and the target of the embedding have the same subsets of κ. Note that if the models have the same subsets of κ, then they have the same objects of transitive size κ from the point of view of the models.

Proposition 2.1. *If κ is a weakly Ramsey cardinal, then κ is a weakly compact limit of weakly compact cardinals.*

Proof. Since κ satisfies the Ramsey embedding property, it has property (4) of Theorem 1.1. So, by Theorem 1.1, to show that κ is weakly compact, it suffices to verify that $\kappa^{<\kappa} = \kappa$. But it is easy to see that κ is inaccessible by using the embeddings to show that it is regular and the κ-powerset preservation property of the embeddings to show that it is a strong limit. Note that without the κ-powerset preservation property it would be impossible to show that κ is a strong limit since property (4) of weakly compact cardinals provably does not imply that κ is a strong limit (see [**?**], ch. 6).

It remains to show that κ is a limit of weakly compact cardinals. We start by choosing a weak κ-model M containing V_κ for which there exists a κ-powerset preserving $j : M \to N$. Observe that it suffices to show that κ is weakly compact in N. If κ is weakly compact in N, then for every $\beta < \kappa$, we have $N \models$ "$\exists \alpha < j(k)\ \alpha > \beta$ and α is weakly compact". It follows that $M \models$ "$\exists \alpha < \kappa\ \alpha > \beta$ and α is weakly compact". But since $V_\kappa \in M$, the model M is correct about α being weakly compact. Thus, it remains to show that $N \models$ "κ is weakly compact". I will show that $N \models$ "Every κ-tree has a cofinal branch". Suppose $T \subseteq \kappa$ is a κ-tree in N. Since M and N have the same subsets of κ, the tree $T \in M$. Consider $j(T) \in N$. Since κ is the critical point of j and levels of T have size less than κ, the tree $j(T)$ restricted to the first κ many levels is exactly T. Since the height of $j(T)$ is $j(\kappa)$, it must have some element on the κ^{th} level. The predecessors of that element will be a cofinal branch of T in N. □

Definition 2.2. An uncountable regular cardinal κ is *ineffable* if for every sequence $\langle A_\alpha \mid \alpha \in \kappa\rangle$ with $A_\alpha \subseteq \alpha$, there exists $A \subseteq \kappa$ such that the set $S = \{\alpha \in \kappa \mid A \cap \alpha = A_\alpha\}$ is stationary. An uncountable regular cardinal κ is *weakly ineffable* if such an A can be found for which the corresponding set S has size κ. A cardinal κ is *ineffably Ramsey* if κ is both ineffable and Ramsey.

Ineffable cardinals are weakly compact limits of weakly compact cardinals and weakly ineffable limits of weakly ineffable cardinals. This is true since ineffable cardinals are Π^1_2-indescribable [**?**] (p. 315), and being weakly ineffable or weakly compact is a Π^1_2- statement satisfied by ineffable cardinals. We will see below that a Ramsey cardinal is a limit of ineffable cardinals. However, since being Ramsey is a Π^1_2-statement, the least Ramsey cardinal cannot be ineffable.

Theorem 2.3. *If κ is a weakly Ramsey cardinal, then κ is a weakly ineffable limit of ineffable cardinals.*

Proof. Fix $\overrightarrow{A} = \langle A_\alpha \mid \alpha \in \kappa\rangle$ with each $A_\alpha \subseteq \alpha$. Choose a weak κ-model M containing $\overrightarrow{A}$ and V_κ for which there exists a κ-powerset preserving embedding $j : M \to N$. Consider $j(\overrightarrow{A})$ and, in particular, $A = j(\overrightarrow{A})(\kappa)$. Since $A \subseteq \kappa$, by the powerset preservation property, $A \in M$. The set $S = \{\alpha \in \kappa \mid A \cap \alpha = A_\alpha\}$ is stationary in M. To see this, fix a club $C \in M$ and observe that $\kappa \in j(S) \cap j(C)$. In particular, S has size κ. So we have shown that κ is weakly ineffable. To show that κ is a limit of ineffable cardinals, we show that $N \models$ "κ is ineffable". The argument above actually shows that κ is ineffable in M. But since M and N have the same subsets of κ, this implies that κ is ineffable in N as well. □

The conclusion of Theorem 2.3 cannot be improved to say that κ is ineffable since a Ramsey cardinal is always weakly Ramsey, and the least Ramsey cardinal is not ineffable. In fact, we show in [**?**] that even the strongly Ramsey cardinals are not necessarily ineffable.

Definition 2.4. An uncountable regular cardinal κ is *subtle* if for every sequence $\langle A_\alpha \mid \alpha \in \kappa\rangle$ with $A_\alpha \subseteq \alpha$ and every club C on κ, there exist ordinals $\alpha < \beta$ in C such that $A_\beta \cap \alpha = A_\alpha$.

Proposition 2.5. *If κ is a weakly Ramsey cardinal, then κ is subtle.*

It is known that weakly ineffable cardinals are subtle, but I will give a direct proof, again using the reflection properties of Ramsey-like embeddings.

Proof. Fix any $\overrightarrow{A} = \langle A_\alpha \mid \alpha \in \kappa\rangle$ with $A_\alpha \subseteq \alpha$ and a club C on κ. Choose a weak κ-model M containing $\overrightarrow{A}$ and C for which there exists a κ-powerset preserving $j : M \to N$. Define S and A as in the proof of Theorem 2.3, and recall that M thinks that S is stationary. Thus, there are $\alpha < \beta$ in $S \cap C$. Clearly for such $\alpha < \beta$, we have $A_\beta \cap \alpha = A \cap \beta \cap \alpha = A_\alpha$. □

Corollary 2.6. *If κ is a weakly Ramsey cardinal, then $\diamondsuit_\kappa$ holds.*

Proof. If κ is subtle, then $\diamondsuit_\kappa$ holds. [3] □

The work of Ian Sharpe [**?**] (see Section 5 for details) shows that:

Theorem 2.7. *If κ is a Ramsey cardinal, then κ is a weakly Ramsey limit of weakly Ramsey cardinals.*

I show in Corollary 3.11 that every strongly Ramsey cardinal is a Ramsey limit of Ramsey cardinals.

Theorem 2.8. *If κ is a super Ramsey cardinal, then κ is a strongly Ramsey limit of strongly Ramsey cardinals.*

Proof. Choose a κ-model $M \prec H_{\kappa^+}$ for which there exists a κ-powerset preserving embedding $j : M \to N$. Note that $V_\kappa \in M$ since M is a κ-model (use the Replacement axiom in M). As usual, it suffices to show that κ is a strongly Ramsey cardinal in N. By Proposition 1.9, there is always a κ-powerset preserving embedding witnessing the strong Ramsey embedding property whose target has size κ. It follows that $H_{\kappa^+} \models$ "κ is a strongly Ramsey cardinal". Therefore $M \models$ "κ is a strongly Ramsey cardinal" by elementarity, and N agrees about this since M and N have the same subsets of κ. □

Theorem 2.9. *If κ is a super Ramsey cardinal, then κ is an ineffably Ramsey limit of ineffably Ramsey cardinals.*

Proof. If κ has the super Ramsey embedding property, then we can choose $M \prec H_{\kappa^+}$ in the proof of Theorem 2.3. This insures that if M thinks that a set is stationary, then M is correct about this. Therefore the proof of Theorem 2.3 shows in this case that κ is ineffable. Once we show that having the Ramsey embedding property is equivalent to being Ramsey (Theorem 3.10), it will follow that κ is also Ramsey. Thus, κ is ineffably Ramsey. We already saw that if $j : M \to N$ is κ-powerset preserving and M is a κ-model, then $N \models$ "κ is ineffable". I will also show later that if

---

[3] See [**?**], p. 315 for a proof that ineffable cardinals have diamond, and observe that only subtleness is needed.

$j : M \to N$ is κ-powerset preserving and M is a κ-model, then $N \models$ "κ is Ramsey" (Corollary 3.11). This shows that κ is a limit of ineffably Ramsey cardinals. $\square$

Corollary 2.10. *A strongly Ramsey cardinal is a limit of ineffably Ramsey cardinals.*

The most surprising result is the following:

Theorem 2.11. *There are no totally Ramsey cardinals.*

Proof. Suppose that there exists a totally Ramsey cardinal and let κ be the *least* totally Ramsey cardinal. Choose any κ-model $M \prec H_{\kappa^+}$ and a κ-powerset preserving embedding $j : M \to N$. The strategy will be to show that κ is totally Ramsey in N. Observe first that $H^N_{\kappa^+} = M$. Thus, to show that κ is totally Ramsey in N, we need to verify in N that every κ-model $m \prec M$ has a κ-powerset preserving embedding. So let $m \in N$ be a κ-model such that $m \prec M$. Observe that $m \in M$ and $m \prec H_{\kappa^+}$. By Proposition 1.9, H_{κ^+} contains a κ-powerset preserving embedding for m. By elementarity, M contains some κ-powerset preserving embedding $h : m \to n$. Since M thinks that $|Trcl(h)| \leq \kappa$, it must be in N as well. Thus, κ is totally Ramsey in N. It follows that there is a totally Ramsey cardinal α below κ. This is, of course, impossible since we assumed that κ was the least totally Ramsey cardinal. Thus, there cannot be any totally Ramsey cardinals. $\square$

This is surprising since, as was pointed out earlier, these embedding properties without powerset preservation are equivalent modulo the assumption that $\kappa^{<\kappa} = \kappa$. Once we add the powerset condition to the embeddings, the equivalence is strongly violated. Of course, now the question arises whether there can be any super Ramsey cardinals.

Theorem 2.12. *If κ is a measurable cardinal, then κ is a super Ramsey limit of super Ramsey cardinals.*

Proof. Fix $A \subseteq \kappa$. Let $j : V \to M$ be an ultrapower by a measure on κ, then j is κ-powerset preserving. First, we reduce j to a κ-powerset preserving embedding of sets by considering the restriction $j : H_{\kappa^+} \to H^M_{j(k)^+}$. The problem is that the set H_{κ^+} is still too big. So second, in some sense, we would like to take an elementary substructure of size κ of the embedding we currently have. To make this precise, consider the structure $\langle H^M_{j(\kappa)^+}, H_{\kappa^+}, j\rangle$. This is a structure whose universe is $H^M_{j(\kappa)^+}$ with a unary relation H_{κ^+} and a binary relation j. Observe that $\langle H^M_{j(\kappa)^+}, H_{\kappa^+}, j\rangle \models$ "j is a κ-powerset preserving embedding from H_{κ^+} to $H^M_{j(\kappa)^+}$". Now we can take an elementary substructure $\langle N', K, h'\rangle$ of size κ such that $A \in N'$, $\kappa + 1 \subseteq N'$, and $N'^{<\kappa} \subseteq N'$. The last is possible, by the usual Skolem-Lowenheim type construction, since κ is inaccessible. By elementarity, we have that $A \in K$, $\kappa + 1 \subseteq K$, $K^{<\kappa} \subseteq K$, and $K \prec H_{\kappa^+}$. Also, since $\kappa \subseteq K$, we have that K is transitive. Thus, K is a κ-model containing A and elementary in H_{κ^+}. Let $\pi : N' \to N$ be the Mostowski collapse and observe that $\pi \restriction K = id$. Finally, let $h = \pi$"h'. Elementarity and the fact that π is an isomorphism imply that $h : K \to N$ is a κ-powerset preserving embedding. This concludes the argument that κ is super Ramsey. It remains to show that κ is a limit of super Ramsey cardinals. It will suffice to show that $M \models$ "κ is super Ramsey". But this follows since the h we built has transitive size κ and is therefore in M. $\square$

Corollary 2.13. *Con*(ZFC$+\exists$ *measurable cardinal*) $\Longrightarrow$ *Con*(ZFC$+\exists$ *proper class of super Ramsey cardinals*)

Next, observe that we cannot hope for any of the cardinals with the Ramsey-like embeddings to have property (8) of weakly compact cardinals from Theorem 1.1. In fact, the following is true:

Proposition 2.14. *If κ has the property that every $A \subseteq \kappa$ is contained in a weak κ-model M for which there exists a κ-powerset preserving embedding $j : M \to N$ such that $j``\mathcal{P}(\kappa)^M$ is an element of N, then κ is a limit of measurable cardinals.*

Proof. Fix a weak κ-model M containing V_κ for which there exists a κ-powerset preserving $j : M \to N$ such that $X = j``\mathcal{P}(\kappa)^M$ is an element of N. Define the usual M-ultrafilter $U = \{B \subseteq \kappa \mid B \in M \text{ and } \kappa \in j(B)\}$. Since $X \in N$, we can define $\{C \cap \kappa \mid C \in X \text{ and } \kappa \in C\} = \{B \subseteq \kappa \mid \kappa \in j(B)\} = U$ in N. Therefore U is an element of N. But, by the powerset preservation property, U is also an N-ultrafilter, and hence N thinks that κ is measurable. It follows that κ must be a limit of measurable cardinals. $\square$

For example, if κ is 2^κ-supercompact, then κ will have the above property. To see this, fix a 2^κ-supercompact embedding $j : V \to M$ and $A \subseteq \kappa$. Choose some cardinal λ such that $j(\lambda) = \lambda$ and $j``2^\kappa \in H^M_{\lambda^+}$. As before, we first restrict j to a set embedding $j : H_{\lambda^+} \to H^M_{\lambda^+}$. Observe that $H^M_{\lambda^+} \subseteq H_{\lambda^+}$. Thus, it makes sense to consider the structure $\langle H_{\lambda^+}, H^M_{\lambda^+}, j\rangle$. Take an elementary substructure $\langle K', N', h'\rangle$ of $\langle H_{\lambda^+}, H^M_{\lambda^+}, j\rangle$ of size κ such that $A \in K'$, $j``2^\kappa \in K'$, $\kappa+1 \subseteq K'$, and $K'^{<\kappa} \subseteq K'$. Let $\pi : K' \to K$ be the Mostowski collapse, $\pi \restriction N' = N$, and $h = \pi``h'$. Observe that N is the Mostowski collapse of N'. It is easy to see that K is a κ-model containing A, the map $h : K \to N$ is a κ-powerset preserving embedding, and $h``\mathcal{P}(\kappa)^K$ is an element of N.

Property (2) of weakly compact cardinals from Theorem 1.1 is called the *Extension Property*. Finally, I will show that the weak Ramsey cardinals also have an extension-like property. Suppose $\mathfrak{X} \subseteq \mathcal{P}(\kappa)$. The structure $\langle V_\kappa, \in, B\rangle_{B\in\mathfrak{X}}$ will be the structure in the language consisting of $\in$ and unary predicate symbols for every element of $\mathfrak{X}$ with the natural interpretation.

Theorem 2.15. *A cardinal κ is weakly Ramsey if and only if every $A \subseteq \kappa$ belongs to a collection $\mathfrak{X} \subseteq \mathcal{P}(\kappa)$ such that the structure $\langle V_\kappa, \in, B\rangle_{B\in\mathfrak{X}}$ has a proper transitive elementary extension $\langle W, \in, B^*\rangle_{B\in\mathfrak{X}}$ with $\mathcal{P}(\kappa)^W = \mathfrak{X}$.*

Proof. ($\Longrightarrow$): Suppose that κ is a weakly Ramsey cardinal and $A \subseteq \kappa$. Fix a weak κ-model M containing A and V_κ for which there exists a κ-powerset preserving embedding $j : M \to N$. Let $\mathfrak{X} = \mathcal{P}(\kappa)^M$. It is easy to verify that $\langle V_\kappa, \in, B\rangle_{B\in\mathfrak{X}} \prec \langle V_{j(\kappa)}, \in, j(B)\rangle_{B\in\mathfrak{X}}$.
($\Longleftarrow$): Fix $A \subseteq \kappa$. The set A belongs to a collection $\mathfrak{X} \subseteq \mathcal{P}(\kappa)$ such that the structure $\langle V_\kappa, \in, B\rangle_{B\in\mathfrak{X}}$ has a proper transitive elementary extension $\langle W, \in, B^*\rangle_{B\in\mathfrak{X}}$ with $\mathcal{P}(\kappa)^W = \mathfrak{X}$. We can assume that W has size κ since if this is not the case, we can take an elementary substructure of size κ which contains V_κ as a subset and collapse it. Since V_κ satisfies that H_{α^+} exists for every $\alpha < \kappa$, it follows by elementarity that H_{κ^+} exists in W. Let $M = H^W_{\kappa^+}$ and observe that M is a weak κ-model containing A. Define $U = \{B \in \mathfrak{X} \mid k \in B^*\}$. I claim that U is a weakly amenable M-ultrafilter. See ahead to Definition 3.2 and Proposition 3.3 for an explanation of weakly amenable ultrafilters. It should be clear that U is

an ultrafilter. To show that U is normal, fix a regressive $f : B \to \kappa$ in M for some $B \in U$. Since we can code f as a subset of κ and $\mathcal{P}(\kappa)^W = \mathfrak{X}$, we can think of f being in $\mathfrak{X}$. Now we can consider the regressive $f^* : B^* \to \kappa^*$ and let $f^*(\kappa) = \alpha < \kappa$. The set $C = \{\xi \in \kappa \mid f(\xi) = \alpha\}$ is in W. Since it is clear that $\kappa \in C^*$, we have $C \in U$. Thus, U is normal. To show that U is weakly amenable, suppose $B \subseteq \kappa \times \kappa$ is in M. We need to see that the set $C = \{\alpha \in \kappa \mid B_\alpha \in U\}$ is in M. Again, since we can code B as a subset of κ, we think of B as being in $\mathfrak{X}$. In W, we can define the set $\{\alpha \in \kappa \mid \kappa \in B^*_\alpha\}$, and it is clear that this set is exactly C. This completes the argument that $C \in M$, and hence U is weakly amenable.

Next, I will show that the ultrapower of M by U is well-founded. It will help first to verify that if $C \in \mathfrak{X}$ codes a well-founded relation on κ, then C^* codes a well-founded relation on Ord^W. If $C \in \mathfrak{X}$ codes a well-founded relation, $\langle V_\kappa, \in, B\rangle_{B \in \mathfrak{X}}$ satisfies that $C \restriction \alpha$ has a rank function for all $\alpha < \kappa$. It follows that $\langle W, \in, B^*\rangle_{B \in \mathfrak{X}}$ satisfies that $C^* \restriction \alpha$ has a rank function for all $\alpha < \text{Ord}^W$. Since κ is weakly compact and we assumed that W has size κ, we can find a *well-founded* elementary extension $\langle X, E, B^{**}\rangle_{B \in \mathfrak{X}}$ for the structure $\langle W, \in, B^*\rangle_{B \in \mathfrak{X}}$ satisfying that there exists an ordinal above the ordinals of W (Theorem 1.1, (1)) . There is no reason to expect that X is an end-extension or that E is the true membership relation, but that is not important for us. We only care that E is well-founded and X thinks it has an ordinal $> \text{Ord}^W$. By elementarity, it follows that $\langle X, E, B^{**}\rangle_{B \in \mathfrak{X}}$ satisfies that $C^{**} \restriction \alpha$ has a rank function for all $\alpha < \text{Ord}^X$. In particular, if $\alpha > \text{Ord}^W$ in X, then $\langle X, E, B^{**}\rangle_{B \in \mathfrak{X}}$ satisfies that $C^{**} \restriction \alpha$ has a rank function. Since the structure $\langle X, E, B^{**}\rangle_{B \in \mathfrak{X}}$ is well-founded and can only add new elements to C^*, if C^* was not well-founded to begin with, X would detect this. Hence C^* is well-founded.

Now we go back to proving that the ultrapower of M by U is well-founded. Suppose towards a contradiction that there exists a membership descending sequence $\ldots E\,[f_n]\,E \ldots E\,[f_1]\,E\,[f_0]$ of elements of the ultrapower. Each $f_n : \kappa \to M$ is an element of M, and for every $n \in \omega$, the set $A_n = \{\alpha \in \kappa \mid f_{n+1}(\alpha) \in f_n(\alpha)\} \in U$. In M, fix some $F_n \subseteq \kappa$ coding the function f_n. We can use the codes F_0 and F_1 and the set A_0 to define $B_0 \subseteq \kappa$, which codes for every $\alpha \in A_0$, a membership isomorphism from the transitive closure of $F_1(\alpha)$ to the subset of the transitive closure of $F_0(\alpha)$ that corresponds to the transitive closure of $f_1(\alpha)$. In this way, we define B_n for every $n \in \omega$. Observe that $\langle V_\kappa, \in, B\rangle_{B \in \mathfrak{X}}$ satisfies that for every $\alpha \in A_0$, the set B_0 codes a membership isomorphism from the transitive closure of $F_1(\alpha)$ to a subset of the transitive closure of $F_0(\alpha)$ that corresponds to the transitive closure of an element of $F_0(\alpha)$. Since $A_0 \in U$, we know that $\kappa \in A_0^*$. Hence $\langle W, \in, B^*\rangle_{B \in \mathfrak{X}}$ satisfies that the set B_0^* codes a membership isomorphism from the transitive closure of $F_1^*(\kappa)$ to the subset of the transitive closure of $F_0^*(\kappa)$ that corresponds to the transitive closure of an element of $F_0^*(\kappa)$. The same statement holds of course for all $n \in \omega$. Since each F_n was well-founded, then so is each F_n^* by the above argument. Hence we can Mostowski collapse each F_n^* to obtain some function $g_n : \text{Ord}^W \to \text{Ord}$. Finally, observe that the $g_n(\kappa)$ form a descending $\in$-sequence. This follows since the transitive closure of each $F_{n+1}^*(\kappa)$ was membership isomorphic to the transitive closure of an element of $F_n^*(\kappa)$. Thus, we reached a contradiction showing that the the ultrapower of M by U is well-founded. Let N be the Mostowski collapse of M/U, and observe finally that since U was weakly amenable, the ultrapower embedding $j : M \to N$ is κ-powerset preserving (Proposition 3.3). □

3. RAMSEY CARDINALS

In this section, I give a proof that Ramsey cardinals are exactly the cardinals with the Ramsey embedding property. Before that we need to recall some facts about iterating M-ultrafilters and κ-powerset preserving embeddings.

Definition 3.1. A cardinal κ is *Ramsey* if every coloring $F:[\kappa]^{<\omega}\to 2$ has a homogenous set of size κ.

Definition 3.2. Suppose M is a weak κ-model and U is an M-ultrafilter on κ. Then U is *weakly amenable* if for every $A\subseteq\kappa\times\kappa$ in M, the set $\{\alpha\in\kappa\mid A_\alpha\in U\}\in M$.

It is easy to see that if U is weakly amenable, then for every $A\subseteq\kappa^n\times\kappa$ in M, the set $\{\vec{\alpha}\in\kappa^n\mid A_{\vec{\alpha}}\in U\}\in M$. If U is a weakly amenable M-ultrafilter on κ, we can define *product* ultrafilters U^n on $\mathcal{P}(k^n)\cap M$ for every $n\in\omega$. We define U^n by induction on n such that $A\subseteq\kappa^n\times\kappa$ is in $U^{n+1}=U^n\times U$ if and only if $A\in M$ and $\{\vec{\alpha}\in\kappa^n\mid A_{\vec{\alpha}}\in U\}\in U^n$. Note that this definition makes sense only in the presence of weak amenability. It turns out that weakly amenable M-ultrafilters on κ are exactly the ones that give rise to κ-powerset preserving embeddings.

Proposition 3.3. *If M is a weak κ-model and $j:M\to N$ is the ultrapower by an M-ultrafilter U on κ, then U is weakly amenable if and only if j is κ-powerset preserving.* ([**?**], p. 246)

Once we have the definition of weakly amenable ultrafilters, we can restate the Ramsey embedding property. We say that an ultrafilter U is *countably closed* if whenever $\langle A_n\mid n\in\omega\rangle$ is a sequence of elements of U, the intersection $\cap_{n\in\omega}A_n\neq\emptyset$.

Proposition 3.4. *A cardinal κ has the Ramsey embedding property if and only if every $A\subseteq\kappa$ is contained in a weak κ-model M for which there exists a countably closed weakly amenable M-ultrafilter on κ.*

This holds since a countably closed ultrafilter must give rise to a well-founded ultrapower. The next couple of propositions address the issue of iterating ultrapowers.

Lemma 3.5. *Suppose M is a weak κ-model, U is a weakly amenable M-ultrafilter on κ, and $j:M\to N$ is the well-founded ultrapower by U^n with critical point κ. Define $j(U)=\{A\subseteq j(\kappa)\mid A=[f]_{U^n}$ and $\{\vec{\alpha}\in\kappa^n\mid f(\vec{\alpha})\in U\}\in U^n\}$. Then $j(U)$ is well-defined and $j(U)$ is a weakly amenable N-ultrafilter on $j(\kappa)$ such that $A\in U$ implies $j(A)\in j(U)$.* ([**?**], p. 246)

Lemma 3.5 is essentially saying that we can take the ultrapower of the structure $\langle M,\in,U\rangle$ by the ultrafilter U^n and the Łoś construction still goes through due to the weak amenability of U. The ultrafilter $j(U)$ is simply the relation corresponding to U in the ultrapower. The next lemma is a modification for the case of weak κ-models of the standard fact from iterating ultrapowers (see, for example, [**?**], ch. 0).

Lemma 3.6. *Suppose M is a weak κ-model, U is a weakly amenable M-ultrafilter on κ, and $j_U:M\to M/U$ is the well-founded ultrapower by U. Suppose further that $j_{U^n}:M\to M/U^n$ and $h_{U^n}:M/U\to(M/U)/U^n$ are the well-founded ultrapowers by U^n. Then the ultrapower $j_{j_{U^n}(U)}:M/U^n\to(M/U^n)/j_{U^n}(U)$ by $j_{U^n}(U)$ and* *the ultrapower* $j_{U^{n+1}} : M \to M/U^{n+1}$ *by* U^{n+1} *are also well-founded. Moreover,* $(M/U^n)/j_{U^n}(U) = (M/U)/U^n = M/U^{n+1}$*, and the following diagram commutes:*

$$\begin{tikzcd}
M \arrow[r, "j_U"] \arrow[d, "j_{U^n}"'] \arrow[dr, "j_{U^{n+1}}"] & M/U \arrow[d, "h_{U^n}"] \\
M/U^n \arrow[r, "j_{j_{U^n}(U)}"'] & (M/U^n)/j_{U^n}(U)
\end{tikzcd}$$

Proposition 3.7. *If* M *is a weak* κ*-model and* U *is a countably closed weakly amenable* M*-ultrafilter on* κ*, then the ultrapowers of* M *by* U^n *are well-founded for all* $n \in \omega$*.*

Proof. Use Lemma 3.6 and argue by induction on n. □

Lemma 3.8. *Suppose* M *is a weak* κ*-model and* U *is a weakly amenable* M*-ultrafilter on* κ *such that the embeddings* $j_{U^n} : M \to M/U^n$ *are well-founded for all* $n \in \omega$*. Then* $A \in U^n$ *if and only if* $\langle \kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)\rangle \in j_{U^n}(A)$*.*

Proof. The proof is by induction on n. Since $j : M \to M/U$ is an ultrapower by a normal ultrafilter, $A \in U$ if and only if $\kappa \in j_U(A)$. This proves the base case, and so assume that the statement holds for n. To prove the statement for $n+1$, we will refer to the commutative diagram from Lemma 3.6. Suppose $\langle \kappa, j_U(\kappa), \ldots, j_{U^n}(\kappa)\rangle \in j_{U^{n+1}}(A)$, then $\langle \kappa, j_U(\kappa), \ldots, j_{U^n}(\kappa)\rangle \in j_{j_{U^n}(U)}(j_{U^n}(A))$, and so $j_{U^n}(\kappa) \in j_{j_{U^n}(U)}(j_{U^n}(A))_{\langle \kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)\rangle}$. It follows that $j_{U^n}(\kappa) \in j_{j_{U^n}(U)}(j_{U^n}(A)_{\langle \kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)\rangle})$ since the critical point of $j_{j_{U^n}(U)}$ is $j_{U^n}(\kappa)$. Since $j_{U^n}(U)$ is a normal ultrafilter on $j_{U^n}(\kappa)$, we have that $j_{U^n}(A)_{\langle \kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)\rangle} \in j_{U^n}(U)$. By the inductive assumption, we know that $[id]_{U^n} = \langle \kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)\rangle$, and therefore we have $j_{U^n}(A)_{[id]_{U^n}} \in j_{U^n}(U)$. But now this implies that $\{\vec{\alpha} \in \kappa^n \mid A_{\vec{\alpha}} \in U\} \in U^n$. So $A \in U^{n+1}$ as desired. For the forward implication, just reverse the steps above. □

Lemma 3.9. *Suppose* M *is a weak* κ*-model and* U *is a weakly amenable* M*-ultrafilter on* κ *such that the ultrapowers by* U^n *are well-founded for all* $n \in \omega$*. If* $A \in U^n$*, then there exists* $B \in U$ *such that for all* $\alpha_1 < \cdots < \alpha_n \in B$*, we have* $\langle \alpha_1, \ldots, \alpha_n\rangle \in A$*.*

Proof. I will argue by induction on n. Throughout the proof, I will be using the machinery of diagonal intersections in disguise, but it is easier to see what is going on without using the terminology of diagonal intersections. The base case $n = 1$ is clearly trivial. So suppose the statement holds for n. Let $A \in U^{n+1}$, then $Z = \{\vec{\alpha} \in \kappa^n \mid A_{\vec{\alpha}} \in U\} \in U^n$. By the inductive assumption, there is a set $Y \in U$ such that for all $\beta_1 < \cdots < \beta_n \in Y$, we have $\langle \beta_1, \ldots, \beta_n\rangle \in Z$. Define $B = \{\xi \in Y \mid \forall \beta_1 < \cdots < \beta_n \in Y \text{if } \beta_n < \xi, \text{ then } \xi \in A_{\langle \beta_1, \ldots, \beta_n\rangle}\}$, then $j(B) =$

$\{\xi \in j(Y) \mid \forall \beta_1 < \cdots < \beta_n \in j(Y) \text{ if } \beta_n < \xi, \text{ then } \xi \in j(A)_{\langle \beta_1,\ldots,\beta_n\rangle}\}$. Recall that $B \in U$ if and only if $\kappa \in j(B)$. It is clear that $\kappa \in j(Y)$. If $\beta_1 < \cdots < \beta_n \in j(Y)$ and $\beta_n < \kappa$, then $\beta_1, \ldots, \beta_n \in Y$. It follows that $\kappa \in j(A_{\langle \beta_1,\ldots,\beta_n\rangle}) = j(A)_{\langle \beta_1,\ldots,\beta_n\rangle}$ since $A_{\langle \beta_1,\ldots,\beta_n\rangle} \in U$. So $\kappa \in j(B)$, and hence $B \in U$. Also, clearly if $\beta_1 < \cdots < \beta_n < \xi$ are in B, then $\langle \beta_1, \ldots, \beta_n, \xi\rangle \in A$. □

We are now ready to prove that cardinals with the Ramsey embedding property are Ramsey. The other direction will involve a much more complicated argument starting with Definition 3.12.

Theorem 3.10. *A cardinal κ has the Ramsey embedding property if and only if κ is Ramsey.*

Proof of the forward direction. Fix a coloring $F : [\kappa]^{<\omega} \to 2$ and choose a weak κ-model M containing F for which there exists a countably closed weakly amenable M-ultrafilter U on κ. Let $j : M \to N$ be the ultrapower by U. We need to show that F has a homogenous set H of size κ. First, observe that if $\langle A_n \mid n \in \omega\rangle$ is a sequence such that each $A_n \in U$, then $\cap_{n\in\omega} A_n$ has size κ. To see this, suppose $\cap_{n\in\omega} A_n$ has size less than κ. Choose $\alpha < \kappa$ such that $\cap_{n\in\omega} A_n$ is contained in α, and let $B = \{\xi \in \kappa \mid \xi > \alpha\} \in U$. But then $B \cap (\cap_{n\in\omega} A_n) = \emptyset$, which contradicts that U is countably closed. Define $f_n : [\kappa]^n \to 2$ by $f_n = F \restriction [\kappa]^n$ and observe that $\langle f_n \mid n \in \omega\rangle \in M$. The strategy will be to find for every $n \in \omega$, a set $H_n \in U$ homogenous for f_n. Then $\cap_{n\in\omega} H_n$ will have size κ and be homogenous for F. Let $j_{U^n}(f_n)(\kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)) = i \in 2$, and consider $A = \{\overrightarrow{\alpha} \in \kappa^n \mid f_n(\overrightarrow{\alpha}) = i\}$. Clearly $A \in U^n$ since $\langle \kappa, j_U(\kappa), \ldots, j_{U^{n-1}}(\kappa)\rangle \in j_{U^n}(A)$. By Proposition 3.9, there exists $B \in U$ such that for all $\beta_1 < \cdots < \beta_n \in B$, we have $\langle \beta_1, \ldots, \beta_n\rangle \in A$. Letting $B = H_n$, we get the desired result. □

Corollary 3.11. *If κ is a strongly Ramsey cardinal, then κ is a Ramsey limit of Ramsey cardinals.*

Proof. Clearly κ is Ramsey. So we need to show that κ is a limit of Ramsey cardinals. Fix a κ-model M for which there exists a κ-powerset preserving $j : M \to N$. The argument above shows that κ is Ramsey in M since M being closed under $<\kappa$-sequences will have $\langle H_n : n \in \omega\rangle$ as an element. Thus, by the powerset preservation property, κ is Ramsey in N as well. Hence, κ is a limit of Ramsey cardinals. □

We now give a summary of the arguments from [**?**] (p. 128, p. 140) that Ramsey cardinals have the Ramsey embedding property.

Definition 3.12. Suppose κ is a cardinal and $A \subseteq \kappa$. Then $I \subseteq \kappa$ is a set of *good indiscernibles* for $\langle L_\kappa[A], A\rangle$ if for all $\gamma \in I$:

(1) $\langle L_\gamma[A], A\rangle \prec \langle L_\kappa[A], A\rangle$.
(2) $I - \gamma$ is a set of indiscernibles for $\langle L_\kappa[A], A, \xi\rangle_{\xi\in\gamma}$.

It turns out that if I are good indiscernibles for $\langle L_\kappa[A], A\rangle$, then every $\gamma \in I$ is inaccessible in $\langle L_\kappa[A], A\rangle$.

Lemma 3.13. *If κ is Ramsey and $A \subseteq \kappa$, then $\langle L_\kappa[A], A\rangle$ has a collection of good indiscernibles of size κ.*

For proof, see [**?**] (p. 140). We are now ready to prove that Ramsey cardinals have the Ramsey embedding property.

Proof of the backward direction of Theorem 3.10. Fix $A \subseteq \kappa$ and consider the structure $\mathcal{A} = \langle L_\kappa[A], A\rangle$. It is clear that $\langle L_\kappa[A], A\rangle$ has a definable well-ordering and therefore definable Skolem functions. By Lemma 3.13, $\langle L_\kappa[A], A\rangle$ has a collection I of good indiscernibles of size κ. For every $\gamma \in I$ and $n \in \omega$, let $\vec{\gamma} = \{\gamma_1, \ldots, \gamma_n\}$ where $\gamma_1 < \cdots < \gamma_n$ are the first n elements in I above γ. Given $\gamma \in I$ and $n \in \omega$, define $\widetilde{\mathcal{A}}^n_\gamma = Scl_{\mathcal{A}}(\gamma + 1 \cup \{\vec{\gamma}\})$, the Skolem closure using the definable Skolem functions of $\langle L_\kappa[A], A\rangle$. It is easy to see that $\langle L_\kappa[A], A\rangle \models \mathrm{ZFC}$. It follows that H_λ exists in $\langle L_\kappa[A], A\rangle$ for every λ, and $\langle H^{\mathcal{A}}_{\lambda^+}, A\rangle \models \mathrm{ZFC}^-$. Since $\langle \widetilde{\mathcal{A}}^n_\gamma, A\rangle \prec \langle L_\kappa[A], A\rangle$ and $\gamma \in \widetilde{\mathcal{A}}^n_\gamma$, we have $H^{\mathcal{A}}_{\gamma^+} \in \widetilde{\mathcal{A}}^n_\gamma$. Next, define $\mathcal{A}^n_\gamma = \widetilde{\mathcal{A}}^n_\gamma \cap H^{\mathcal{A}}_{\gamma^+}$.

Lemma 3.13.1. *$\mathcal{A}^n_\gamma$ is transitive and $\langle \mathcal{A}^n_\gamma, A\rangle \models \mathrm{ZFC}^-$.*

Proof. First, we show that $\mathcal{A}^n_\gamma$ is transitive. Fix $a \in \mathcal{A}^n_\gamma$ and $b \in a$. The set a is coded by a subset of $\gamma \times \gamma$ in $\langle L_\kappa[A], A\rangle$. By elementarity, $\widetilde{\mathcal{A}}^n_\gamma$ contains a set $E \subseteq \gamma \times \gamma$ coding a and the Mostowski collapse $\pi : \langle \gamma, E\rangle \to Trcl(a)$. Let $\alpha \in \gamma$ such that $\langle L_\kappa[A], A\rangle \models \pi(\alpha) = b$. Since $\alpha < \gamma$, we have $\alpha \in \widetilde{\mathcal{A}}^n_\gamma$, and so $b \in \widetilde{\mathcal{A}}^n_\gamma$ by elementarity. It is clear that $b \in \mathcal{A}^n_\gamma$ also. This shows that $\mathcal{A}^n_\gamma$ is transitive.

Next, we show that $\langle \mathcal{A}^n_\gamma, A\rangle \models \mathrm{ZFC}^-$. Let $\pi : \widetilde{\mathcal{A}}^n_\gamma \to N$ be the Mostowski collapse. Since $\mathcal{A}^n_\gamma \subseteq \widetilde{\mathcal{A}}^n_\gamma$ is transitive, it follows that $\pi(x) = x$ for all $x \in \mathcal{A}^n_\gamma$. Therefore $\pi(H_{\gamma^+}) = \mathcal{A}^n_\gamma$ and $\pi(H_{\gamma^+} \cap A) = \mathcal{A}^n_\gamma \cap A$. It follows that $\langle \mathcal{A}^n_\gamma, A\rangle$ is an element of the Mostowski collapse N and N thinks $\mathcal{A}^n_\gamma$ is its H_{γ^+}. From this we conclude that $\langle \mathcal{A}^n_\gamma, A\rangle \models \mathrm{ZFC}^-$. $\square$

Lemma 3.13.2. *For every $\gamma \in I$ and $n \in \omega$, we have $\mathcal{A}^n_\gamma \prec \mathcal{A}^{n+1}_\gamma$.*

Proof. It is certainly clear that $\widetilde{\mathcal{A}}^n_\gamma \prec \widetilde{\mathcal{A}}^{n+1}_\gamma$ since these are Skolem closures of sets that extend each other. Let $\pi : \widetilde{\mathcal{A}}^n_\gamma \to N$ and $\rho : \widetilde{\mathcal{A}}^{n+1}_\gamma \to N'$ be the Mostowski collapses. It suffices to observe that $\mathcal{A}^n_\gamma$ is the H_{γ^+} of N and $\mathcal{A}^{n+1}_\gamma$ is the H_{γ^+} of N'. $\square$

Recall that if $a \in \widetilde{\mathcal{A}}^n_\gamma$, then $a = h(\vec{b}, \gamma, \vec{\gamma})$ where h is a definable Skolem function, $\vec{b} \subseteq \gamma$, and $\vec{\gamma} = \{\gamma_1, \ldots, \gamma_n\}$ are the first n elements above γ in I. Given $\gamma < \delta \in I$, define $f^n_{\gamma\delta} : \widetilde{\mathcal{A}}^n_\gamma \to \widetilde{\mathcal{A}}^n_\delta$ by $f^n_{\gamma\delta}(a) = h(\vec{b}, \delta, \vec{\delta})$ where $a = h(\vec{b}, \gamma, \vec{\gamma})$ is as above. Observe that since $I - \gamma$ are indiscernibles for $\langle L_\kappa[A], A, \xi\rangle_{\xi \in \gamma}$, the map $f^n_{\gamma\delta}$ is clearly well-defined and elementary. Also, $f^n_{\gamma\delta}(\gamma) = \delta$ and $f^n_{\gamma\delta}(\xi) = \xi$ for all $\xi < \gamma$. So the critical point of $f^n_{\gamma\delta}$ is γ. Finally, note that for all $\gamma < \delta < \beta \in I$, we have $f^n_{\gamma\beta} \circ f^n_{\beta\delta} = f^n_{\gamma\delta}$.

Lemma 3.13.3. *The map $f^n_{\gamma\delta} \restriction \mathcal{A}^n_\gamma : \mathcal{A}^n_\gamma \to \mathcal{A}^n_\delta$ is elementary.*

Proof. Fix $a \in \mathcal{A}^n_\gamma$ and recall that $\widetilde{\mathcal{A}}^n_\gamma$ thinks $a \in H_{\gamma^+}$. By elementarity of $f^n_{\gamma\delta}$, it follows that $\widetilde{\mathcal{A}}^n_\delta$ thinks $f^n_{\gamma\delta}(a) \in H_{\delta^+}$. Therefore $f^n_{\gamma\delta} : \mathcal{A}^n_\gamma \to \mathcal{A}^n_\delta$. Elementarity follows as in the previous lemma. $\square$

For $\gamma \in I$, define $U^n_\gamma = \{X \in \mathcal{P}(\gamma) \cap \mathcal{A}^n_\gamma \mid \gamma \in f^n_{\gamma\delta}(X) \text{ for some } \delta > \gamma\}$. Equivalently, we could have used "for *all* $\delta > \gamma$" in the definition.

Lemma 3.13.4. *U^n_γ is an $\mathcal{A}^n_\gamma$-ultrafilter on γ.*

Proof. Easy. $\square$

Observe that if $a_0, \ldots, a_n \in L_\gamma[A]$ for some $\gamma \in I$, then for every formula $\varphi(\overrightarrow{x})$, we have $\langle L_\kappa[A], A\rangle \models \varphi(\overrightarrow{a}) \leftrightarrow \langle L_\gamma[A], A\rangle \models \varphi(\overrightarrow{a}) \leftrightarrow \langle L_\kappa[A], A\rangle \models$ "$\langle L_\gamma[A], A\rangle \models \varphi(\overrightarrow{a})$". It follows that for every $\gamma \in I$, the model $\langle L_\kappa[A], A\rangle$ has a truth predicate definable from γ for formulas with parameters from $L_\gamma[A]$.

Lemma 3.13.5. *The ultrafilter $U^n_\gamma \in \mathcal{A}^{n+2}_\gamma$.*

Proof. In $\widetilde{\mathcal{A}}^{n+2}_\gamma$, we have $U^n_\gamma = \{x \in \mathcal{P}(\gamma) \mid \exists h\, \exists \overrightarrow{v} < \gamma\ h \text{ is a Skolem term and } h = (\overrightarrow{v}, \gamma, \gamma_1, \ldots, \gamma_n) \wedge \gamma \in h(\overrightarrow{v}, \gamma_1, \gamma_2, \ldots, \gamma_{n+1})\}$. This follows since $\gamma_{n+2} \in \widetilde{\mathcal{A}}^{n+2}_\gamma$, and therefore we can define a truth predicate for $L_{\gamma_{n+2}}[A]$, which is good enough for the definition above. So far we have shown that U^n_γ is in $\widetilde{\mathcal{A}}^{n+2}_\gamma$. To finish the argument, observe that $\langle L_\kappa[A], A\rangle \models U^n_\gamma \in H_{\gamma^+}$, and therefore $U^n_\gamma \in \mathcal{A}^{n+2}_\gamma$. $\square$

It should be clear that $U^n_\gamma \subseteq U^{n+1}_\gamma$ and $f^n_{\gamma\delta} \subseteq f^{n+1}_{\gamma\delta}$. Define $\mathcal{A}_\gamma = \cup_{n\in\omega} \mathcal{A}^n_\gamma$ and $U_\gamma = \cup_{n\in\omega} U^n_\gamma$. Also, define $f_{\gamma\delta} = \cup_{n\in\omega} f^n_{\gamma\delta} : \mathcal{A}_\gamma \to \mathcal{A}_\delta$ and observe that it is elementary. Since each U^n_γ was an $\mathcal{A}^n_\gamma$-ultrafilter on γ, it is easy to see that U_γ is an $\mathcal{A}_\gamma$-ultrafilter on γ.

Lemma 3.13.6. *The $\mathcal{A}_\gamma$-ultrafilter U_γ is weakly amenable.*

Proof. Consider $B \subseteq \gamma \times \gamma$ with $B \in \mathcal{A}_\gamma$. Define $C = \{\xi \in \gamma \mid B_\xi \in U_\gamma\}$. We need to show that $C \in \mathcal{A}_\gamma$. Since $B \in \mathcal{A}_\gamma$, it follows that $B \in \mathcal{A}^n_\gamma$ for some $n \in \omega$. But then $C = \{\xi \in \gamma \mid B_\xi \in U^n_\gamma\}$ and $U^n_\gamma \in \mathcal{A}^{n+2}_\gamma \subseteq \mathcal{A}_\gamma$. $\square$

Now for every $\gamma \in I$, we have an associated structure $\langle \mathcal{A}_\gamma, \in, A, U_\gamma\rangle$. Also, if $\gamma < \delta$ in I, we have an elementary embedding $f_{\gamma\delta} : \mathcal{A}_\gamma \to \mathcal{A}_\delta$ with critical point γ between the structures $\langle \mathcal{A}_\gamma, A\rangle$ and $\langle \mathcal{A}_\delta, A\rangle$ such that $X \in U_\gamma$ if and only if $f_{\gamma\delta}(X) \in U_\delta$. This is a directed system of models and so we can take its direct limit. Define $\langle B, E, A', V\rangle = \lim_{\gamma\in I} \langle \mathcal{A}_\gamma, \in, A, U_\gamma\rangle$.

Lemma 3.13.7. *The relation E on B is well-founded.*

Proof. The elements of B are functions t with domains $\{\xi \in I \mid \xi \geq \alpha\}$ for some $\alpha \in I$ satisfying the properties:

1. $t(\gamma) \in \mathcal{A}_\gamma$,
2. for $\gamma < \delta$ in domain of t, we have $t(\delta) = f_{\gamma\delta}(t(\gamma))$,
3. there is no $\xi \in I \cap \alpha$ for which there is $a \in \mathcal{A}_\xi$ such that $f_{\xi\alpha}(a) = t(a)$.

Note that each t is determined once you know any $t(\xi)$ by extending uniquely forward and backward. Standard arguments (for example, [**?**], p. 157) show that $B \models \varphi(t_1, \ldots, t_n) \leftrightarrow \exists \gamma\, \mathcal{A}_\gamma \models \varphi(t_1(\gamma), \ldots, t_n(\gamma)) \leftrightarrow$ for all γ in the intersection of the domains of the t_i, the structure $\mathcal{A}_\gamma \models \varphi(t_1(\gamma), \ldots, t_n(\gamma))$. This truth definition holds only of atomic formulas where the formulas involve the predicate for the ultrafilter.

Suppose to the contrary that E is not well-founded, then there is a descending E-sequence $\cdots E\, t_n\, E \cdots E\, t_1\, E\, t_0$. Find γ_0 such that $\mathcal{A}_{\gamma_0} \models t_1(\gamma_0) \in t_0(\gamma_0)$. Next, find $\gamma_1 > \gamma_0$ such that $\mathcal{A}_{\gamma_1} \models t_2(\gamma_1) \in t_1(\gamma_1)$. In this fashion, define an increasing sequence $\gamma_0 < \gamma_1 < \cdots < \gamma_n < \cdots$ such that $\mathcal{A}_{\gamma_n} \models t_{n+1}(\gamma_n) \in t_n(\gamma_n)$.

Let $\gamma \in I$ such that $\gamma > \sup_{n\in\omega}\gamma_n$. It follows that for all $n \in \omega$, the structure $\mathcal{A}_\gamma \models f_{\gamma_n\gamma}(t_{n+1}(\gamma_n)) \in f_{\gamma_n\gamma}(t_n(\gamma_n))$, and therefore $\mathcal{A}_\gamma \models t_{n+1}(\gamma) \in t_n(\gamma)$. But, of course, this is impossible. Thus, E is well-founded. $\square$

Let $\langle \mathcal{A}_\kappa, \in, A^*, U_\kappa\rangle$ be the Mostowski collapse of $\langle B, E, A', V\rangle$.

Lemma 3.13.8. *The cardinal* $\kappa \in \mathcal{A}_\kappa$.

Proof. Fix $\alpha \in \kappa$ and let $\gamma \in I$ be the least ordinal greater than α. Define t_α to have domain $\{\xi \in I \mid \xi \geq \gamma\}$ such that $t_\alpha(\xi) = \alpha$, then t_α is an element of B which collapses to α. Define t_κ to have domain I such that $t_\kappa(\gamma) = \gamma$, then t_κ collapses to κ. $\square$

Define $j_\gamma : \mathcal{A}_\gamma \to \mathcal{A}_\kappa$ such that $j_\gamma(a)$ is the collapse of the function t for which $t(\gamma) = a$. The maps j_γ are fully elementary in the language of $\langle A_\gamma, A\rangle$ and elementary for atomic formulas in the language with the predicate for the ultrafilter. Observe that $j_\gamma(\xi) = \xi$ for all $\xi < \gamma$ since if $t(\gamma) = \xi$, then $t = t_\xi$. Also, $j_\gamma(\gamma) = \kappa$ since if $t(\gamma) = \gamma$, then $t = t_\kappa$. So the critical point of each j_γ is κ. Finally, if $\gamma < \delta$ in I, then $j_\delta \circ f_{\gamma\delta} = j_\gamma$.

Lemma 3.13.9. *The set* U_κ *is a weakly amenable* $\mathcal{A}_\kappa$*-ultrafilter on* κ.

Proof. Easy. $\square$

Lemma 3.13.10. *A set* $X \in U_\kappa$ *if and only if there exists* $\alpha \in I$ *such that* $\{\xi \in I \mid \xi > \alpha\} \subseteq X$.

Proof. Fix $X \subseteq \kappa$ in $\mathcal{A}_\kappa$ and $\beta \in I$ such that for all $\xi > \beta$, there is $X' \in \mathcal{A}_\xi$ with $j_\xi(X') = X$. For $\xi > \beta$, we have $X \in U_\kappa \leftrightarrow X' \in U_\xi \leftrightarrow \xi \in f_{\xi\xi_1}(X') \leftrightarrow j_{\xi_1}(\xi) \in j_{\xi_1} \circ f_{\xi\xi_1}(X') = j_\xi(X') \leftrightarrow \xi \in j_\xi(X') = X$. Thus, for $\alpha > \beta$, we have $\{\xi \in I \mid \xi > \alpha\} \subseteq X$ if and only if $X \in U_\kappa$. $\square$

Lemma 3.13.11. *The* $\mathcal{A}_\kappa$*-ultrafilter* U_κ *is countably closed.*

Proof. Fix $\langle A_n \mid n \in \omega\rangle$ a sequence of elements of U_κ. We need to show that $\cap_{n\in\omega} A_n \neq \emptyset$. For each A_n, there exists $\gamma_n \in I$ such that $X_n = \{\xi \in I \mid \xi > \gamma_n\} \subseteq A_n$. Thus, $\cap_{n\in\omega} X_n \subseteq \cap_{n\in\omega} A_n$ and clearly $\cap_{n\in\omega} X_n$ has size κ. $\square$

It remains to show that $A \in \mathcal{A}_\kappa$.

Lemma 3.13.12. *The set* $A^* \restriction \kappa = A$*, and hence* $A \in \mathcal{A}_\kappa$.

Proof. Fix $\alpha \in A$ and let $\gamma \in I$ such that $\gamma > \alpha$, then $\langle \mathcal{A}_\gamma, A\rangle \models \alpha \in A$. It follows that $\langle \mathcal{A}_\kappa, A\rangle \models j_\gamma(\alpha) \in A^*$, but $j_\gamma(\alpha) = \alpha$, and so $\alpha \in A^*$. Thus, $A \subseteq A^*$. Now fix $\alpha \in A^* \restriction \kappa$ and let $\gamma \in I$ such that $\gamma > \alpha$, then $j_\gamma(\alpha) = \alpha$, and so $j_\gamma(\alpha) \in A^*$. It follows that $\alpha \in A$. Thus, $A^* \restriction \kappa \subseteq A$. We conclude that $A = A^* \restriction \kappa$. $\square$

We have found a weak κ-model M containing A for which there exists a countably closed weakly amenable M-ultrafilter on κ, namely $M = \mathcal{A}_\kappa$. This concludes the proof that if κ is Ramsey, then κ has the Ramsey embedding property. $\square$

I will end this section by giving an interesting reformulation of the Ramsey embedding property.

Proposition 3.14. *A cardinal* κ *is Ramsey if and only if every* $A \subseteq \kappa$ *is contained in a weak* κ*-model* $M \models \mathrm{ZFC}$ *for which there exists* $j : M \to N$ *an ultrapower by a countably closed* M*-ultrafilter such that* $M \prec N$.

The difference from the earlier definition of the Ramsey embedding property is that now for every $A \subseteq \kappa$, we have $A \in M$ where M is a model of *all of* ZFC and $j : M \to N$ is an ultrapower by a countably closed M-ultrafilter such that not only do M and N have the same subsets of κ, but actually $M \prec N$.

Proof. Fix $A \subseteq \kappa$ and choose a weak κ-model M containing A and V_κ for which there exists a countably closed weakly amenable M-ultrafilter U on κ. Let $j : M \to N$ be the ultrapower by U. I will refer to the commutative diagram from Lemma 3.6. For the case $n = 1$, the diagram becomes the following:

$$\begin{tikzcd}
M \arrow[r, "j"] \arrow[d, "j"'] \arrow[dr, "j_{U_2}"] & N = M/U \arrow[d, "h_U"] \\
N = M/U \arrow[r, "j_{j(U)}"'] & K = N/j(U)
\end{tikzcd}$$

Let $M' = V^N_{j(\kappa)}$, then M' is a transitive model of ZFC since $V_\kappa \models$ ZFC. Let $K' = V^K_{j_{j(U)}(j(\kappa))} = V^K_{h_U(j(\kappa))}$. Observe that since M' is transitive, the map $h_U \restriction M' : M' \to K'$ is the ultrapower embedding of M' into M'/U. It remains to show that $M' \prec K'$, but this follows easily from the $j_{j(U)}$ side of the commutative diagram since $j_{j(U)} \restriction M' : M' \to N'$ and $j_{j(U)}$ is identity on M'. □

Corollary 3.15. *A cardinal κ is strongly Ramsey if and only if every $A \subseteq \kappa$ is contained in a κ-model $M \models$ ZFC for which there exists an elementary embedding $j : M \to N$ with critical point κ such that $M \prec N$.*

Proof. Using the previous proof it suffices to show that M' is closed under $< \kappa$-sequences. We can assume without loss of generality that N is closed under $< \kappa$-sequences (Proposition 1.9). Therefore $V^N_{j(\kappa)}$ must be closed under $< \kappa$-sequences since N thinks that $j(\kappa)$ is inaccessible. □

Question 3.16. If κ is a weakly Ramsey cardinal, does there exist for every $A \subseteq \kappa$, a weak κ-model $M \models$ ZFC and an elementary embedding $j : M \to N$ with critical point κ such that $M \prec N$?

In the above proofs, we get elementary embeddings $j : M \to N$ such that $M = V^N_{j(\kappa)}$. It is worth noting that it is always possible to get κ-powerset preserving embeddings where $M = H^N_{\kappa^+}$.

Proposition 3.17. *If $j : M \to N$ is a κ-powerset preserving embedding of weak κ-models and $A \subseteq \kappa$ is in M, then there is another κ-powerset preserving embedding of weak κ-models $h : M' \to N'$ such that $A \in M'$ and $M' = H^{N'}_{\kappa^+}$.*

Proof. Define $M' = \{x \in M \mid M \models |Trcl(x)| \leq \kappa\}$ and $N' = \{x \in N \mid N \models |Trcl(x)| \leq j(\kappa)\}$. Clearly M' is a transitive subclass of M containing A. It is easy to check that $j \restriction M' : M' \to N'$ is a κ-powerset preserving embedding of κ-models and $M' = H^{N'}_{\kappa^+}$. $\square$

4. Weakly Ramsey Cardinals Exist in L

In this section, I show that weakly Ramsey cardinals are downward absolute to L. The proof is nearly completely due to Philip Welch who showed originally that if $0^\#$ exists, then Silver indiscernibles are weakly Ramsey in L [?]. I modified the proof slightly using Lemma 4.3 to obtain the full downward absoluteness.

Lemma 4.1. *If $0^\#$ exists and κ is a Silver indiscernible, then $(\kappa^+)^L$ has cofinality ω.*

Proof. Let I be the Silver indiscernibles and let $\kappa \in I$. Every element of L has the form $t(\overrightarrow{\beta}, \overrightarrow{\gamma})$ where t is a Skolem term and $\overrightarrow{\beta} \leq \kappa$ and $\overrightarrow{\gamma} > \kappa$ are both from I. Fix a Skolem term t and let $k = m + n$ be the arity of t. Let $\delta_1 < \cdots < \delta_n$ be the first n indiscernibles above κ. We will investigate all possible values of the term t with parameters from I such that the first m many parameters are below or equal to κ and the n remaining parameters are above κ. I claim that for $\overrightarrow{\beta} \leq \kappa$ and $\overrightarrow{\gamma} > \kappa$ in I, if $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) < (\kappa^+)^L$, then $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) = t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n)$. Since $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) < (\kappa^+)^L$, by indiscernibility we have that $t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n) < (\kappa^+)^L$. It follows that for any n many indiscernibles $\xi_1 < \cdots < \xi_n$ above κ, we have $t(\overrightarrow{\beta}, \xi_1, \ldots, \xi_n) < (\kappa^+)^L$. Split the indiscernibles above κ into class many increasing n-tuples. That is, we have $\delta_1 < \cdots < \delta_n < \delta_{n+1} < \cdots < \delta_{2n} < \cdots$. If $t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n) < t(\overrightarrow{\beta}, \delta_{n+1}, \ldots, \delta_{2n})$, then, by indiscernibility, we have class many elements below κ^+, which is impossible. Also, we can't have $t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n) > t(\overrightarrow{\beta}, \delta_{n+1}, \ldots, \delta_{2n})$ since, by indiscernibility, we would have a descending ω-chain in the ordinals. It follows that we must have $t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n) = t(\overrightarrow{\beta}, \delta_{n+1}, \ldots, \delta_{2n})$. Now let $\gamma_n, \delta_n < \delta_{\xi_1} < \cdots < \delta_{\xi_n}$, then $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) = t(\overrightarrow{\beta}, \delta_{\xi_1}, \ldots, \delta_{\xi_n})$ and $t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n) = t(\overrightarrow{\beta}, \delta_{\xi_1}, \ldots, \delta_{\xi_n})$. This gives $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) = t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n)$. Thus, for a fixed term t of arity $k = m + n$ and $\overrightarrow{\beta} \leq \kappa$ of arity m in I, we have for all $\overrightarrow{\gamma} > \kappa$ of arity n in I that if $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) < (\kappa^+)^L$, then $t(\overrightarrow{\beta}, \overrightarrow{\gamma}) = t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n)$ where $\overrightarrow{\delta}$ is as above. Fix such t and consider the supremum in L of $t(\overrightarrow{\beta}, \delta_1, \ldots, \delta_n) < (\kappa^+)^L$ ranging over all $\overrightarrow{\beta} < \kappa$ and $n \leq k$. Since this supremum is definable in L and L thinks that $(\kappa^+)^L$ is regular, it follows that the supremum is below $(\kappa^+)^L$. Thus, $(\kappa^+)^L$ is the union of these supremums taken over all terms t, and hence has cofinality ω. $\square$

Theorem 4.2 (Welch, 2007). *If $0^\#$ exists, then Silver indiscernibles are weakly Ramsey in L.*

Proof. Let $I = \{i_\xi : \xi \in \mathrm{Ord}\}$ be the Silver indiscernibles enumerated in order. Fix $\kappa \in I$ and let $\lambda = (\kappa^+)^L$. Let $j : I \to I$ such that $j(i_\xi) = i_\xi$ for all $i_\xi < \kappa$ and $j(i_\xi) = i_{\xi+1}$ for all $i_\xi \geq \kappa$ in I. The map j generates, via the Skolem functions, the elementary embedding $j : L \to L$ with critical point κ. We can restrict j to $j : L_\lambda \to L_{j(\lambda)}$. Since $\lambda = (\kappa^+)^L$, it should be clear that j is κ-powerset preserving. Let U be the weakly amenable L_λ-ultrafilter generated by κ via j (Proposition 1.9). Observe that, by weak amenability, $U_\alpha = U \cap L_\alpha$ is an element of L_λ for all $\alpha < \lambda$ since α has size κ in L_λ. Construct, using Lemma 4.1, a sequence $\langle\langle L_{\lambda_i}, U_{\lambda_i}\rangle : i \in \omega\rangle$ such that $\langle \lambda_i : i \in \omega\rangle$ is cofinal in λ, each $L_{\lambda_i} \prec L_\lambda$, and $U_{\lambda_i}, L_{\lambda_i} \in L_{\lambda_{i+1}}$. Note that each individual $\langle L_{\lambda_i}, U_{\lambda_i}\rangle$ is an element of L and each U_{λ_i} is a L_{λ_i}-ultrafilter with a *well-founded* ultrapower, but not necessarily weakly amenable. This motivates the argument below.

Fix $A \subseteq \kappa$ in L. Define in L, the tree T of finite sequences of the form $s = \langle\langle L_{\gamma_0}, W_0, \varphi_0\rangle, \ldots, \langle L_{\gamma_n}, W_n, \varphi_n\rangle\rangle$ ordered by extension and satisfying the properties:

1. $A \in L_{\gamma_0}$,
2. $L_{\gamma_i} \prec L_\lambda$ and W_i is an L_{γ_i}-ultrafilter on κ with a *well-founded* ultrapower,
3. φ_i is a map from the equivalence classes in $Ult(L_{\gamma_i}, W_i)$ which represent ordinals order-preserving into Ord,
4. for $i < j \leq n$, we have $L_{\gamma_i}, W_i \in L_{\gamma_j}$, $W_i \subseteq W_j$, and "$\varphi_i \subseteq \varphi_j$".

Since $L_{\gamma_i} \prec L_{\gamma_j}$ and $W_i \subseteq W_j$, we have that $Ult(L_{\gamma_i}, W_i)$ elementarily embeds into $Ult(L_{\gamma_j}, W_j)$ via the canonical map $[f]_{W_i} \mapsto [f]_{W_j}$. The quotation marks around $\varphi_i \subseteq \varphi_j$ above refer to the inclusion modulo to this embedding.

The sequence $\langle\langle L_{\lambda_i}, U_{\lambda_i}\rangle : i \in \omega\rangle$ constructed above gives a branch through the tree T in V since the maps φ_{λ_i} can be easily constructed from the canonical embeddings of $Ult(L_{\lambda_i}, U_{\lambda_i})$ into $L_{j(\lambda)}$. Thus, the tree is ill-founded in V, and hence also ill-founded in L by absoluteness. Let $\{\langle L_{\gamma_i}, W_i, \varphi_i\rangle : i \in \omega\}$ be a branch of T in L. Let $L_\gamma = \cup_{i\in\omega} L_{\gamma_i}$, $W = \cup_{i\in\omega} W_i$, and $\varphi = \cup_{i\in\omega}\varphi_i$. It is clear that W is a *weakly amenable* L_γ-ultrafilter. The function φ is an order-preserving map from the ordinals of $Ult(L_\gamma, W)$ into Ord witnessing that the ultrapower by W is *well-founded*. To see this, fix $[f]_W$ and $[g]_W$ representing ordinals in $Ult(L_\gamma, W)$. Since $L_\gamma = \cup_{i\in\omega} L_{\gamma_i}$, there is $\gamma_i < \gamma$ such that f and g are elements of L_{γ_i}. Thus, $[f]_{W_i}$ and $[g]_{W_i}$ represent ordinals in $Ult(L_{\gamma_i}, W_i)$ and the function φ_i mapped them order-preserving into Ord. We now found a weak κ-model L_γ containg A for which there exists a weakly amenable L_γ-ultrafilter on κ with a well-founded ultrapower. This completes the proof that κ is weakly Ramsey. □

Next, I will prove a lemma that will allow us to modify the proof of Theorem 4.2 to show that weakly Ramsey cardinals are downward absolute to L.

Lemma 4.3. *If κ is a weakly Ramsey cardinal, then every $A \subseteq \kappa$ is contained in a weak κ-model M of cofinality ω for which there exists a κ-powerset preserving embedding $j : M \to N$.*

Proof. Fix $A \subseteq \kappa$ and choose a κ-powerset preserving embedding $h : M' \to N'$ such that $A \in M'$ and $M' = H^{N'}_{\kappa^+}$ (Proposition 3.17). I claim that if $B \in M'$, then $h \restriction B \in N'$. In M', let f be a bijection from κ onto B and consider $h(f) \in N'$. Fix $b \in B$ and let $f(\xi) = b$, then $h(f)(\xi) = h(f)(h(\xi)) = h(f(\xi)) = h(b)$. Thus $h \restriction B = h(f)f^{-1}$. Define $X'_0 = \kappa + 1 \cup \{A\}$, then X'_0 is an element of size κ in N'. Let $X_0 = X'_0 \cup h``(X'_0 \cap M')$, then X_0 is an element of size κ in N'. Note that subsets of H_{κ^+} of size κ are also elements of H_{κ^+}, and hence $X'_0 \cap M'$ is an element of M'. Define X'_1 to be the closure of X_0 under witnesses for Σ_1-formulas with parameters from X_0 satisfied by N'. Again X'_1 is an element of size κ in N'. Let $X_1 = X'_1 \cup h``(X'_1 \cap M')$. Similarly, define $X'_{i+1}$ to be the closure of $X_i$ under witnesses for $\Sigma_{i+1}$-formulas with parameters from $X_i$ satisfied by $N'$ and let $X_{i+1} = X'_{i+1} \cup h``(X'_{i+1} \cap M')$. Define $X = \cup_{i\in\omega} X_i$. It is clear that $X \prec N'$. Let $\pi : X \to N$ be the Mostowski collapse and let $M = H^N_{\kappa^+}$. It should be clear that N and M both have cofinality ω. Define $j : M \to N$ by $j(a) = \pi(h(\pi^{-1}(a)))$. The definition makes sense since if $a \in M$, we have $\pi^{-1}(a) \in M'$, and hence $h(\pi^{-1}(a)) \in X$ by construction. It is clear that h is κ-powerset preserving. So it suffices to verify that h is elementary. Suppose $M \models \varphi(a)$, then $X \models$ "$H_{\kappa^+} \models \varphi(\pi^{-1}(a))$", and hence $N' \models$ "$H_{\kappa^+} \models \varphi(\pi^{-1}(a))$". It follows that $M' \models \varphi(\pi^{-1}(a))$, and hence $N' \models \varphi(h(\pi^{-1}(a)))$. This implies that $X \models \varphi(h(\pi^{-1}(a)))$, and hence $N \models \varphi(\pi(j(\pi^{-1}(a))))$. $\square$

Theorem 4.4. *If κ is weakly Ramsey, then κ is weakly Ramsey in L.*

Proof. Suppose κ is weakly Ramsey and fix $A \subseteq \kappa$ in L. In L, let μ be an ordinal of size κ such that $A \in L_\mu$ and $B \subseteq \kappa$ a code for L_μ. In V, let M be a weak κ-model of *cofinality* ω containing B for which there exists a κ-powerset preserving $j : M \to N$ (Lemma 4.3). Suppose also that M has the least possible height. Note that the procedure to build M with cofinality ω from the proof of Lemma 4.3 can only decrease the height, so the assumptions are not contradictory. Let $\alpha = \text{Ord}^M$ and $\beta = \text{Ord}^N$, then $L^M = L_\alpha$ and $L^N = L_\beta$. Consider the restriction of j to $j : L_\alpha \to L_\beta$. I will argue that it is still κ-powerset preserving. By running the usual condensation argument inside L_β, it suffices to show that $(\kappa^+)^{L_\beta} \leq \alpha$. Let $\gamma < \kappa^+$ in L_β, then L_β has $C \subseteq \kappa$ coding γ. This C is an element of M, and hence $\gamma \in M$. Thus, $\gamma < \alpha$. Next, I claim that $(\kappa^+)^{L_\beta} = \alpha$. If not, then there is $\xi < \alpha$ such that L_α thinks ξ is κ^+. But then $B \in L_\xi$ and the restriction $j : L_\xi \to L_{j(\xi)}$ is κ-powerset preserving, contradicting that we chose M of least height. Therefore, L_α thinks that all $\gamma < \alpha$ have size κ. Let U be the weakly amenable L_α-ultrafilter generated by κ via j (Proposition 1.9). Observe that, by weak amenability, $U_\gamma = U \cap L_\gamma$ is an element of L_α for all $\gamma < \alpha$ since L_α thinks all $\gamma < \alpha$ have size κ. Now use the fact that α has cofinality ω to proceed as in the proof of Theorem 4.2. $\square$

5. NEARLY RAMSEY CARDINALS

One of the central features of M-ultrafilters that arise from Ramsey-like embeddings is that they can be iterated. For a weak κ-model M, say that an M-ultrafilter on κ is *1-good* if the ultrapower is well-founded. Recall that a cardinal κ is weakly compact if $\kappa^{<\kappa} = \kappa$ and every $A \subseteq \kappa$ is a contained in a weak κ-model for which there exists a 1-good M-ultrafilter on κ. Once we assume that the M-ultrafilters are also weakly amenable, we get that the cardinal is already weakly Ramsey, a much stronger large cardinal. The weak amenablity of an ultrafilter makes possible the notion of iterating it.

Start with a weak κ-model M_0 for which there exists a weakly amenable 1-good M_0-ultrafilter U_0. Applying Lemma 3.5, take the ultrapower of the structure $\langle M_0, U_0\rangle$ by U_0 to obtain the structure $\langle M_1, U_1\rangle$ where U_1 is a weakly amenable M_1-ultrafilter. This is a *one-step* iteration. If the ultrapower of M_1 by U_1 happens to be well-founded, we will say that the original ultrafilter U_0 is *2-good* and take the ultrapower of $\langle M_1, U_1\rangle$ by U_1 to obtain the structure $\langle M_2, U_2\rangle$. This is a *two-step* iteration. Proceeding in this manner, suppose that an M_0-ultrafilter U_0 is *n-good* for every $n \in \omega$. Then we say that U_0 is *ω-good.* In this case, we have a directed (linear actually) system of models $\langle M_n, U_n\rangle$ with the corresponding ultrapower embeddings. Take the direct limit of this system. If the direct limit happens to be well-founded, collapse it to obtain $\langle M_\omega, U_\omega\rangle$, where U_ω is a weakly amenable

M_ω-ultrafilter, and say that the original ultrafilter U_0 is $\omega+1$-*good*. Proceed in this fashion as long as the iterates are well-founded. See [**?**], p. 244 for details involved in this construction. Call an ultrafilter *iterable* if it is α-good for every α. Gaifman showed in [**?**] that if an ultrafilter is ω_1-good, then it is already iterable . Kunen showed in [**?**] that if an ultrafilter is countably complete, then it is ω_1-good, and hence iterable. Thus, every M-ultrafilter that arises from a Ramsey embedding is iterable.

Definition 5.1. A cardinal κ is *nearly Ramsey* if every $A \subseteq \kappa$ is contained in a weak κ-model M for which there exists an iterable M-ultrafilter on κ.

Ramsey cardinals are nearly Ramsey and nearly Ramsey cardinals imply $0^{\#}$. It suffices to show that there is an embedding $j : L_{\kappa^+} \to L_\xi$ with critical point κ. Choose any weak κ-model M for which there exists an iterable M-ultrafilter on κ and consider the iterate M_α whose ordinals include $(\kappa^+)^L$. Then L_{κ^+} is contained in M_α, but M and M_α have the same subsets of κ. Thus, L_{κ^+} is already contained in M, and hence we can restrict the embedding to L_{κ^+}. Most of the work on nearly Ramsey cardinals has been done by Ian Sharpe [**?**]. He showed that nearly Ramsey cardinals are strictly weaker than Ramsey cardinals. In fact, he showed that ω_1-Erdős cardinals are limits of nearly Ramsey cardinals, and hence Ramsey cardinals are limits of nearly Ramsey cardinals. Among other things, this trivially implies that Ramsey cardinals are limits of weakly Ramsey cardinals since every nearly Ramsey is weakly Ramsey.

Question 5.2. Do α-good cardinals form a hierarchy of strength?

Question 5.3. For what ordinals α are α-good cardinals consistent with $V = L$?

Question 5.4. Is every nearly Ramsey cardinal a limit of weakly Ramsey cardinals?

New York City College of Technology (CUNY), 300 Jay Street, Brooklyn, NY 11201
USA
E-mail address: `vgitman@nylogic.org`